

ON SOME SYSTEMS OF DIFFERENCE EQUATIONS

L.A.Gutnik

*Dedicated to the memory of
Professor N.M.Korobov.*

Let

$$|z| \geq 1, -3\pi/2 < \arg(z) \leq \pi/2, \log(z) = \ln(|z|) + i \arg(z).$$

Then $\log(-z) = \log(z) - i\pi$, if $\Re(z) > 0$ and $\log(z) = \log(-z) - i\pi$, if $\Re(z) < 0$.

Let

$$(1) \quad f_{l,1}(z, \nu) = \sum_{k=0}^{\nu} (-1)^{(\nu+k)l} (z)^k \binom{\nu}{k}^{2+l} \binom{\nu+k}{\nu}^{2+l},$$

where $l = 0, 1, 2$, $\nu \in [0, +\infty) \cap \mathbb{Z}$. Let

$$(2) \quad R(t, \nu) = \frac{\prod_{j=1}^{\nu} (t-j)}{\prod_{j=0}^{\nu} (t+j)}.$$

where $\nu \in [0, +\infty) \cap \mathbb{Z}$,

$$(3) \quad f_{l,2}(z, \nu) = \sum_{t=1+\nu}^{+\infty} z^{-t} (R(t, \nu))^{2+l},$$

where $l = 0, 1, 2$ and $\nu \in [0, +\infty) \cap \mathbb{Z}$, and since $(R(t, \nu))^{2+l}$ for $\nu \in \mathbb{N}$ has in the points $t = 1, \dots, \nu$, the zeros of the order $2+l$, it follows that

$$(4) \quad f_{l,2}(z, \nu) = \sum_{t=1}^{+\infty} z^{-t} (R(t, \nu))^{2+l},$$

for $l = 0, 1, 2$ and $\nu \in [0, +\infty) \cap \mathbb{Z}$. Let

$$(5) \quad f_{l,3}(z, \nu) = (\log(z)) f_{l,2}(z, \nu) + f_{l,4}(z, \nu),$$

where

$$(6) \quad f_{l,4}(z, \nu) = - \sum_{t=1+\nu}^{+\infty} z^{-t} \left(\frac{\partial}{\partial t} (R^{2+l}) \right) (t, \nu),$$

$l = 0, 1, 2$ and $\nu \in [0, +\infty) \cap \mathbb{Z}$, and since $(R(t, \nu))^{2+l}$ for $\nu \in \mathbb{N}$ has in the points $t = 1, \dots, \nu$, the zeros of the order $2 + l$, it follows that

$$(7) \quad f_{l,4}(z, \nu) = - \sum_{t=1}^{+\infty} z^{-t} \left(\frac{\partial}{\partial t} (R^{2+l}) \right) (t, \nu)$$

for $l = 0, 1, 2$ and $\nu \in [0, +\infty) \cap \mathbb{Z}$. Let

$$(8) \quad f_{l,5}^{\vee}(z, \nu) = -i\pi f_{l,3}(z, \nu) + f_{l,5}(z, \nu),$$

with $l = 1, 2, \nu \in [0, +\infty) \cap \mathbb{Z}$ and

$$(9) \quad f_{l,5}(z, \nu) = \\ 2^{-1}(\log(z))^2 f_{l,2}(z, \nu) + (\log(z)) f_{l,4}(z, \nu) + f_{l,6}(z, \nu) = \\ = -2^{-1}(\log(z))^2 f_{l,2}(z, \nu) + (\log(z)) f_{l,3}(z, \nu) + f_{l,6}(z, \nu),$$

where

$$(10) \quad f_{l,6}(z, \nu) = 2^{-1} \sum_{t=1+\nu}^{\infty} z^{-t} \left(\left(\frac{\partial}{\partial t} \right)^2 (R^{2+l}) \right) (t, \nu),$$

and since $(R(t, \nu))^{2+l}$ for $\nu \in \mathbb{N}$ has in the points $t = 1, \dots, \nu$, the zeros of the order $2 + l$, and $l = 1, 2$ now, it follows that

$$(11) \quad f_{l,6}(z, \nu) = 2^{-1} \sum_{t=1+\nu}^{\infty} z^{-t} \left(\left(\frac{\partial}{\partial t} \right)^2 (R^{2+l}) \right) (t, \nu)$$

for $l = 1, 2$ and $\nu \in [0, +\infty) \cap \mathbb{Z}$. Let

$$(12) \quad f_{l,7}^{\vee}(z, \nu) = f_{l,7}(z, \nu) + (2\pi^2/3) f_{l,3}(z, \nu).$$

with $l = 2, \nu \in [0, +\infty) \cap \mathbb{Z}$ and

$$(13) \quad f_{l,7}(z, \nu) = \\ -3^{-1}(\log(z))^3 f_{l,2}(z, \nu) + 2^{-1}(\log(z))^2 f_{l,3}(z, \nu) + f_{l,8}(z, \nu) + \\ (\log(z))(f_{l,5}(z, \nu) + 2^{-1}(\log(z))^2 f_{l,2}(z, \nu) - (\log(z)) f_{l,3}(z, \nu)) = \\ 6^{-1}(\log(z))^3 f_{l,2}(z, \nu) - 2^{-1}(\log(z))^2 f_{l,3}(z, \nu) + (\log(z)) f_{l,5}(z, \nu) + f_{l,8}(z, \nu),$$

where

$$(14) \quad f_{l,8}(z, \nu) = -6^{-1} \sum_{t=\nu+1}^{\infty} z^{-t} \left(\left(\frac{\partial}{\partial t} \right)^3 (R^{2+l}) \right) (t, \nu),$$

and, since $(R(t, \nu))^{2+l}$ for $\nu \in \mathbb{N}$ have in the points $t = 1, \dots, \nu$, the zeros of the order $2 + l$, and $l = 2$ now, it follows that

Let

$$\mathfrak{K}_0 = \{1, 2, 3\}, \mathfrak{K}_1 = \{1, 2, 3, 5\}, \mathfrak{K}_2 = \{1, 2, 3, 5, 7\}.$$

Let λ be a variable. We denote by $T_{n,\lambda}$ the diagonal $n \times n$ -matrix, i -th diagonal element of which is equal to λ^{i-1} for $i = 1, \dots, n$. We denote by δ the operator $z \frac{d}{dz}$. Let further $l = 0, 1, 2, k \in \mathfrak{K}_l, |z| > 1, \nu \in \mathbb{N}$, and let $Y_{l,k}(z; \nu)$ be the column with $4 + 2l$ elements, i -th of which is equal to $(\nu^{-1}\delta)^{i-1} f_{l,k}^\vee(z, \nu)$ for $i = 1, \dots, 4 + 2l$.

Theorem 1. *The following equalities hold*

$$(16) \quad A_l^\sim(z; \nu) Y_{l,k}(z; \nu) = T_{4+2l, 1-\nu^{-1}} Y_{l,k}(z; \nu - 1),$$

$$(17) \quad Y_{l,k}(z; \nu) = T_{4+2l, -1} A_l^\sim(z; -\nu) T_{4+2l, -1+\nu^{-1}} Y_{l,k}(z; \nu - 1),$$

where $l = 0, 1, 2, k \in \mathfrak{K}_l, |z| > 1, \nu \in \mathbb{N}, \nu \geq 2$,

$$A_l^\sim(z; \nu) = S_l^\sim + z \sum_{i=0}^{1+l} \nu^{-i} V_l^{\sim*}(i)$$

with

$$(18) \quad S_0^\sim = \begin{pmatrix} 1 & -4 & 8 & -12 \\ 0 & 1 & -4 & 8 \\ 0 & 0 & 1 & -4 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$(19) \quad S_1^\sim = \begin{pmatrix} -1 & 6 & -18 & 38 & -66 & 102 \\ 0 & -1 & 6 & -18 & 38 & -66 \\ 0 & 0 & -1 & 6 & -18 & 38 \\ 0 & 0 & 0 & -1 & 6 & -18 \\ 0 & 0 & 0 & 0 & -1 & 6 \\ 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix},$$

$$(20) \quad S_2^\sim = \begin{pmatrix} 1 & -8 & 32 & -88 & 192 & -360 & 608 & -952 \\ 0 & 1 & -8 & 32 & -88 & 192 & -360 & 608 \\ 0 & 0 & 1 & -8 & 32 & -88 & 192 & -360 \\ 0 & 0 & 0 & 1 & -8 & 32 & -88 & 192 \\ 0 & 0 & 0 & 0 & 1 & -8 & 32 & -88 \\ 0 & 0 & 0 & 0 & 0 & 1 & -8 & 32 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -8 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$V_0^{\sim*}(0) = 4 \begin{pmatrix} 4 & -5 & -2 & 3 \\ -3 & 4 & 1 & -2 \\ 2 & -3 & 0 & 1 \\ -1 & 2 & -1 & 0 \end{pmatrix},$$

$$V_0^{\sim*}(1) = 4 \begin{pmatrix} 3 & -6 & 3 & 0 \\ -2 & 4 & -2 & 0 \\ 1 & -2 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$V_1^{\sim*}(0) = \begin{pmatrix} 146 & -198 & -180 & 268 & 66 & -102 \\ -102 & 146 & 108 & -180 & -38 & 66 \\ 66 & -102 & -52 & 108 & 18 & -38 \\ -38 & 66 & 12 & -52 & -6 & 18 \\ 18 & -38 & 12 & 12 & 2 & -6 \\ -6 & 18 & -20 & 12 & -6 & 2 \end{pmatrix},$$

$$V_1^{\sim*}(1) = \begin{pmatrix} 240 & -516 & 108 & 372 & -204 & 0 \\ -160 & 348 & -84 & -236 & 132 & 0 \\ 96 & -212 & 60 & 132 & -76 & 0 \\ -48 & 108 & -36 & -60 & 36 & 0 \\ 16 & -36 & 12 & 20 & -12 & 0 \\ 0 & -4 & 12 & -12 & 4 & 0 \end{pmatrix},$$

$$V_1^{\sim*}(2) = \begin{pmatrix} 102 & -306 & 306 & -102 & 0 & 0 \\ -66 & 198 & -198 & 66 & 0 & 0 \\ 38 & -114 & 114 & -38 & 0 & 0 \\ -18 & 54 & -54 & 18 & 0 & 0 \\ 6 & -18 & 18 & -6 & 0 & 0 \\ -2 & 6 & -6 & 2 & 0 & 0 \end{pmatrix},$$

$$V_2^{\sim*}(0) = 8 \begin{pmatrix} 176 & -249 & -364 & 545 & 280 & -431 & -76 & 119 \\ -119 & 176 & 227 & -364 & -169 & 280 & 45 & -76 \\ 76 & -119 & -128 & 227 & 92 & -169 & -24 & 45 \\ -45 & 76 & 61 & -128 & -43 & 92 & 11 & -24 \\ 24 & -45 & -20 & 61 & 16 & -43 & -4 & 11 \\ -11 & 24 & -1 & -20 & -5 & 16 & 1 & -4 \\ 4 & -11 & 8 & -1 & 4 & -5 & 0 & 1 \\ -1 & 4 & -7 & 8 & -7 & 4 & -1 & 0 \end{pmatrix},$$

$$V_2^{\sim*}(1) = 8 \begin{pmatrix} 455 & -1020 & -113 & 1552 & -603 & -628 & 357 & 0 \\ -300 & 682 & 44 & -996 & 404 & 394 & -228 & 0 \\ 185 & -428 & -3 & 592 & -253 & -228 & 135 & 0 \\ -104 & 246 & -16 & -316 & 144 & 118 & -72 & 0 \\ 51 & -124 & 19 & 144 & -71 & -52 & 33 & 0 \\ -20 & 50 & -12 & -52 & 28 & 18 & -12 & 0 \\ 5 & -12 & 1 & 16 & -9 & -4 & 3 & 0 \\ 0 & -2 & 8 & -12 & 8 & -2 & 0 & 0 \end{pmatrix},$$

$$V_2^{\sim*}(2) = 8 \begin{pmatrix} 400 & -1243 & 972 & 542 & -1028 & 357 & 0 & 0 \\ -259 & 808 & -642 & -332 & 653 & -228 & 0 & 0 \\ 156 & -489 & 396 & 186 & -384 & 135 & 0 & 0 \\ -85 & 268 & -222 & -92 & 203 & -72 & 0 & 0 \\ 40 & -127 & 108 & 38 & -92 & 33 & 0 & 0 \\ -15 & 48 & -42 & -12 & 33 & -12 & 0 & 0 \\ 4 & -13 & 12 & 2 & -8 & 3 & 0 & 0 \\ -1 & 4 & -6 & 4 & -1 & 0 & 0 & 0 \end{pmatrix},$$

$$V_2^{\sim*}(3) = 8 \begin{pmatrix} 119 & -476 & 714 & -476 & 119 & 0 & 0 & 0 \\ -76 & 304 & -456 & 304 & -76 & 0 & 0 & 0 \\ 45 & -180 & 270 & -180 & 45 & 0 & 0 & 0 \\ -24 & 96 & -144 & 96 & -24 & 0 & 0 & 0 \\ 11 & -44 & 66 & -44 & 11 & 0 & 0 & 0 \\ -4 & 16 & -24 & 16 & -4 & 0 & 0 & 0 \\ 1 & -4 & 6 & -4 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

The above matrices $S_l^{\sim}$ and $V_l^{\sim*}(i)$ have the following properties:

$$(21) \quad A_l^{\sim}(z; -\nu) T_{4+2l, -1} A_{l, k}(z; \nu) = T_{4+2l, -1},$$

$$(22) \quad S_l^{\sim} T_{4+2l, -1} = (S_l^{\sim} T_{4+2l, -1})^{-1}$$

$$(23) \quad S_l^{\sim} T_{4+2l, -1} V_l^{\sim*}(i) = -(-1)^i V_l^{\sim*}(i) T_{4+2l, -1} S_l^{\sim},$$

$$(24) \quad V_l^{\sim*}(i) T_{4+2l, -1} V_l^{\sim*}(k) = 0 T_{4+2l, -1},$$

where

$$l = 0, 1, 2, i \in [0, 1 + l] \cap \mathbb{Z}, k \in [0, 1 + l] \cap \mathbb{Z}.$$

Proof. Full proof can be found in [2] – [6].

References.

- [1] R.Apéry, Interpolation des fractions continues
et irrationalite de certaines constantes,
Bulletin de la section des sciences du C.T.H., 1981, No 3, 37 – 53;
- [2] L.A.Gutnik, On some systems of difference equations. Part 1.
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2006, 23, 1 – 37.
- [3] ———, On some systems of difference equations. Part 2.
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2006, 49, 1 – 31.
- [4] ———, On some systems of difference equations. Part 3.
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2006, 91, 1 – 52.
- [5] ———, On some systems of difference equations. Part 4.
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2006, 101, 1 – 49.
- [6] ———, On some systems of difference equations. Part 5.
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2006, 115, 1 – 9.

E-mail: gutnik@gutnik.mccme.ru